

May 11, 2016

filename bibk.doc

- Kácha, P. and V. Petr. 1996. Camouflage and mimicry in fossils, I.: General part. -- Acta musei Nationalis Pragae, Series B, Historia Naturalis 51(1995)(1-4):53-82. [a very stimulating paper] [p. 58 graptolite colonies imitating brittlestar per Archie Lamont][p. 67 strategy of being different -- example of Ophiopholis aculeata (see paper by G.B. Moment in fh files)] [p. 67 Ordovician predation]
- Kammer, T. W., and W. I. Ausich. 2007. Soft-tissue preservation of the hind gut in a new genus of cladid crinoid from the Mississippian (Visean, Asbian) at St. Andrews, Scotland. Palaeontology 50(4):951-959. [cites soft-tissue preservation in Bundenbachia beneckeii, Palaeophiomyxa grandis, and Protasterina flexuosa; cites Glass 2006; also Glass & Blake 2002, 2004] [this locality is in Fife, Scotland, so perhaps this paper relates also to Fife Aganaster]
- Katzer, Friedrich. 1902. Geologie von Bohmen. Zweite Ausgabe. Prag. Verlag von I. Taussig. [On p. 870 mentions Asterias primula Barr.]
- Kayser, E. 1897. Beiträge zur Kenntnis einiger paläozoischer Faunen Süd-Amerikas. Zeitschrift der Deutschen Geologischen Gesellschaft 49:274-317, pl. 7-12. [see Haude 1999] [listed Reich 2004]
- Kelly, J. 1860. On the Graywacke rocks of Ireland, as compared with those of England. Journal of the Geological Society of Dublin 8:251-333. [p. 318 lists of fossils including Bdellacoma] [Echinodermata on pp. 317-318]
- Kelly, S. M. 1984. Fauna and paleoecology of the Middle Chester Indian Springs Shale, Sulphur, Indiana. Unpub. PhD dissertation, Indiana University, Bloomington. [source Blake & Elliott 2001 ms for JP; new specimen of Delicaster enigmaticus IU#16226]
- Kent, L. S. 1982. Type and figured fossils in the Worthen collection at the Illinois State Geological Survey. Illinois Department of Energy and Natural Resources, State Geological Survey Division, Circular 524, 65 pp. [Protaster? gregarius p. 52; Cholaster peculiaris p. 26; Tremataster difficilis p. 62; Palasterina (Schoenaster) fimbriata p. 45; Compsaster formosus p. 27]
- Kesling, R. V. 1962. Notes on Protopalaeaster narrawayi Hudson. Jour. Paleont., vol. 36, no. 5, pp. 933-942, illus., table.
- Kesling, R. V. 1964. A drastic reappraisal of "Lepidasterella babcocki Schuchert" as Helianthaster gyalinus Clarke a streptophiuran auluroid. Contr. Mus. Pal. Univ. Mich. vol. XIX, no. 9, pp. 115-133, 4 pls., 1 fig.

May 11, 2016

- Kesling, R. V. 1967. Neopalaeaster enigmaticus, new starfish from Upper Mississippian Paint Creek Formation in Illinois. Mich. Univ. Mus. Pal. Contr. vol. 21, no. 3.
- Kesling, R. V. 1969. Silicaster, a new genus of Devonian starfish. Contr. Mus. Pal. Univ. Mich. vol. 22, no. 19, pp. 249-261, 4 pls., 2 text-figs.
- Kesling, Robert V. 1969. Three Permian starfish from Western Australia and their bearing on revision of the Asteroidea. Contr. Mus. Pal. Univ. Mich., vol. 22, no. 25, pp. 361-376, 3 pls., 6 text-figs.
- Kesling, Robert V. 1969. A new brittle-star from the Middle Devonian Arkona Shale of Ontario. Contr. Mus. Pal. Univ. Mich., vol. 23, no. 2, pp. 37-51, 6 pls., 2 text-figs.
- Kesling, Robert V. 1970. Drepanaster wrighti, a new species of brittle-star from the Middle Devonian Arkona Shale of Ontario. Contr. Mus. Pal. Univ. Mich., vol. 23, no. 4, pp. 73-79, 2 pls.
- Kesling, Robert V. 1971. Antiquaster magrumi, a new unusual brittle-star from the Middle Devonian Silica Formation of Northwestern Ohio. Contr. Mus. Pal. Univ. Mich., vol. 23, no. 10, pp. 181-191, 4 pls., 1 text-fig.
- Kesling, Robert V. 1971. Michiganaster inexpectatus, a new many-armed starfish from the Middle Devonian Rogers City limestone of Michigan. Contrib. Mus. Paleont. Univ. Mich., vol. 23, no. 16, pp. 247-262, 3 pls., 5 text-figs.
- Kesling, Robert V. 1972. Strataster devonicus, a new brittle-star with unusual preservation from the Middle Devonian Silica Formation of Ohio. Contr. Mus. Pal. Univ. Mich., vol. 24, no. 2, pp. 9-15, 2 pls., 3 text-figs.
- Kesling, Robert V. 1982. Acinetaster konieckii, a new brittle-star from the Middle Devonian Arkona Shale. Contrib. Mus. Paleont. Univ. Mich. 26(5):71-81.
- Kesling, Robert V. 1982. Arkonaster, a new multiradiate starfish from the Middle Devonian Arkona Shale of Ontario. Contrib. Mus. Paleont. Univ. Mich. 26(6):83-115. [p. 98 table and key to known Middle Devonian Stelleroidea from Michigan, Ohio and Ontario]
- Kesling, Robert V. 1993. Asteroids. Geological Survey of Western Australia, Bulletin, 136: 68, pl. 62. [source Petr]
- Kesling, Robert V. and Ruth B. Chilman. 1975. Strata and megafossils of the Middle Devonian Silica Formation. Friends of the University of Michigan Museum of Paleontology, Inc. Papers on Paleontology No. 8, 408 pp. [Antiquaster magrumi; ?Devonaster spp.,

May 11, 2016

‘multiarmed starfish’ Silicaster esseri, Strataster devonicus] [8/24/2003 Mike Topor: compare Arkonaster with the multiarmed starfish – do they appear to be the same genus/species?] [p. 169 tiny flat oral spines on the lateral plates; millions of these little flat spatula-shaped spines in the shale...]

Kesling, Robert V. and Ruth B. Chilman. 1978. Ostracods of the Middle Devonian Silica Formation. Museum of Paleontology, The University of Michigan, Papers on Paleontology No. 18: vol. 1 (text) & vol. 2 (plates). [Bursulella records; see Hotchkiss & Glass 2010]

Kesling, Robert V. and Douglas Le Vasseur. 1971. Strataster ohioensis, a new Early Mississippian brittle-star, and the paleoecology of its community. Contr. Mus. Pal. Univ. Mich., vol. 23, no. 20, pp. 305-341, 9 text-figs, 13 plates.

Kesling, R. V. and H. L. Strimple. 1966. Calliasterella americana, a new starfish from the Pennsylvanian of Illinois. Jour. Paleont., vol. 40, no. 5, pp. 1157-1166, illus. [they use the convention that the madreporite marks the right anterior interradius; an indentation in the central plate suggests the anus was right posterior]

Kesling, R. V. and Jean D. Wright. 1965. Two new Middle Devonian species of Devonaster from Southwestern Ontario. Mich. Univ. Mus. Pal. Contr., vol. 20, no 15, pp. 257-263.

Kettnerová, M. 1933. Fossilní hvězdice z porýnského devonu. Věda přírodní (Praha), 14(2/3): 35-39. [source Petr]

Keyes, Charles Rollin and B. H. Beane. 1934. Modernity in Paleozoic starfishes. Pan-Am. Geologist, vol. 62, no. 3, pp. 197-212, 2 figs, 3 pls.

Keyes, Richard. 2015. Fossils of the Tennessee Valley. Fossil identification guide for the Tennessee Valley. 131 pp. [Asterozoa pp. 94-95]

Kier, P. M. 1973. The echinoderms and Permian-Triassic time. Pp. 622-629 In The Permian and Triassic Systems and their mutual boundary. Canadian Society of Petroleum Geologists, Memoir 2. 766pp.

Kilfoyle, Clinton F. 1954. Catalog of type specimens of fossils in the New York State Museum, supplement 4. NYSM Bull. 348, 719 pp. [Asterozoa on pp. 196-204, 619; also in listing by strata.]

Kilfoyle, Clinton F. 1969. Catalog of type specimens of fossils in the New York State Museum, supplement 6. NYSM Bull 413, 305 pp. [Devonaster eucharis p. 258, 293.]

Kilius, R. 1969. Stamm Echinodermata - Stachelhäuter. In: Urania Tierreich: Wirbellose Tiere 2 (Annelida bis Chaetognatha), 505-591, Urania-Verlag. Leipzig, Jena, Berlin.

May 11, 2016

Kinahan, G. H. 1878. [see Baily 1878 in Kinahan 1878]

Kinahan, G. H. and W. H. Baily. 1879. Explanatory Memoir to accompany Sheets 169, 170, 180 and 181 of the Map of the Geological Survey of Ireland in the County of Wexford with Palaeontological notes by W. H. Baily. *Memoirs of the Geological Survey of Ireland*. 63 pp. [ref to p. 59 fig. 6] [see Parkes & Sleeman 1997]

Kindle, E. M. 1898. A catalog of the fossils of Indiana: accompanied by a bibliography of the literature relating to them. pp. 407-514 *in* Indiana Department of Geology and Natural Resources, twenty-second annual report [for] 1897. 1197pp.

Kirk, Edwin. 1942. Ampelocrinus, a new crinoid genus from the Upper Mississippian.--*American Journal of Science* 240:22-28. [Plate 1 fig. 2 of paratype of A. bernhardinae from Glen Dean limestone (Chester Group), Sloans Valley, Pulaski County, Ky. has evident ophiuroid cf. Encrinasteridae, number S4402b in the Springer collection of the USNM]

Klages, O. 1955. Die versteckten Fossilien von Bundenbach. *Aufschluss* 6:66-68, 6 figs. Rossdorf bei Darmstadt. [source Kutscher 1976] [Loriolaster mirabilis, Aspidosoma tischbeinianum] [not seen]

Klähn, H. 1929. Die Bedeutung der Seelilien und Seesterne für die Erkennung von Wasserbewegung nach Richtung und Stärke. *Palaeobiologica* 2:287-302, plates 26-34. Wien & Leipzig. [source Kutscher 1976] [Furcaster palaeozoicus, Aspidosoma tischbeinianum, Asterias asperula]

Klinghardt, F. 1933. Beobachtung an lebenden und fossilen Schlangen-Seesternen, Quallen und Rudisten. *Jahrbuch der Preussischen Geologischen Landesanstalt zu Berlin für das Jahr 1932*, vol. 53, pp. 947-963, 7 figs., plates 49-53, Berlin, 1933. [Roemeraster asperula, Furcaster palaeozoicus, Cheiropteraster giganteus, Loriolaster mirabilis] [also Mesozoic and Recent ophiuroids] [arm enrollment in O. fürstenbergii and mention of Onychaster flexilis] [O. fürstenbergii is tentatively Ophioplinthaca fide Jagt (2000 *Scripta Geologica* 121, p. 11)]

Kneidl, V. 1984. Hunsrück und Nahe: Geologie, Mineralogie und Paläontologie; ein Wegweiser für den Liebhaber. Kosmos Gesellschaft der Naturfreunde, Franckh'sche Verlagshandlung. Stuttgart. 128 pp. [faunal list of Asterozoa; photos of Eospondylus, Furcaster, Hymenosoma, Palaeosolaster, Loriolaster]

Knod, Reinhold. 1908. Beiträge zur Geologie und Palaontologie von Sudamerika. XIV. Devonische Faunen Boliviens. *Neues Jahrbuch für Mineralogie, Geologie und Palaontologie*. XXV. Beilage-Band. pp. 493-600. [Records starfish remains from

May 11, 2016

"Pflanzensandstein von Challa".]

Köhler, S. 2002. Im Tode vereint. Fossilien 19(6):323 [Bohemura jahni] [Letna-Park, Prague, 1997]

Kokorin, Alexandr I. 2012. [Russian language abstract] p. 26 Modern paleontology: classical and newest methods. The ninth all-Russian scientific school for young scientists in paleontology. October 1-3, 2012. Borissiak Paleontological Institute of the Russian Academy of Sciences, Moscow. Abstracts. Tezisu_9_shkola.pdf [Ophiuroidea; *Stenaster confluens*]

Kokorin, Alexandr. 2014. Brittle stars in the Carboniferous of the Moscow Basin. EEC Abstracts, Plymouth, UK. no pagination. [*Stenaster confluens* an ophiuroid; new material; also microfossils of Ophiurida and of Furcaster]

Kokorin, Alexandr. 2015. What can *Aganaster* vertebrae tell about its ecology? pp. 83-85 In S. Zamora & I. Rábano, eds., Progress in echinoderm palaeobiology. Cuadernos del Museo Geominero, 19, Instituto Geológico y Minero de España, Madrid ISBN 978-84-7840-961-7. [new material, articulated and also isolated remains; Upper Carboniferous, Moscow Basin] [SEM of isolated vertebra described, interpreted] [comparisons made with *Ophiotholia* and *Ophiomyces*]

Kolata, Dennis R. 1975. Middle Ordovician echinoderms from Northern Illinois and Southern Wisconsin. Paleont. Soc., Mem. 7 (Jour. Paleontology, vol. 49, no. 3, supp.), 74 pp. [Hudsonaster sp. cf. H. rugosus (Billings) on p. 54; Promopalaeaster wilsoni (Raymond) on p. 54; Urasterella? sp. on p. 55] [plate 10 figs 1-4, 7-9]

Kolata, D. R., J. C. Brower & T. J. Frest. 1987. Upper Mississippi Valley Champlainian and Cincinnati echinoderms. pp. 179-181 in R. E. Sloan (ed.) Middle and Late Ordovician lithostratigraphy and biostratigraphy of the upper Mississippi Valley. Minnesota Geological Survey Report of Investigations 35. 232 pp. [University of Minnesota] [Hudsonaster, Promopalaeaster, Schuchertia, Urasterella]

Kolata, D. R. and L. Pavlides. 1986. Echinoderms from the Arvonian slate, central Virginia piedmont.--Geologica et Palaeontologica 20:1-9, illustr. [ZR1987/88] [p.4 ophiuroid (order undetermined)] [see Watson & Powell 1911] [Middle or Upper Ordovician]

Königswald, R. von. 1930. Die Arten der Einregelung im Sediment bei den Seesternen und Seelilien des unterdevonischen Bundenbacher Schiefers. Senckenbergiana, 12:338-360, 19 figs., 7 pls. Frankfurt. [eleven gen. sp. figured] [source Kutscher 1976]

Königswald, R. von. 1930. Die Fauna des Bundenbacher Schiefers in ihren Beziehungen zum Sediment. Cbl. Miner. B:241-247. Stuttgart. [source Kutscher 1976] [not seen]

May 11, 2016

- Kooser, M. 1995. Drawings by Marilyn Kooser, University of California, Riverside p. 490 In: J.D. Cooper, M.L. Droser & S.C. Finney (eds.): *Ordovician Odyssey: Short papers for the Seventh International Symposium on the Ordovician System*, Las Vegas, Nevada, USA, June 1995, 498 pp., The Pacific Section Society for Sedimentary Geology (SEPM). Fullerton, California. [drawing of *Hudsonaster*] [these drawings correspond with illustrations in paper by M.L. Droser, R.A. Fortey & X. Li 1996]
- Kornicker, Louis S. 1954. Distribution of ostracodes in the Florena Shale. Master's Essay, Columbia University, 85 pp. [Columbia University, Butler Library, Microform Reading Room (on microform; no call number available)] [Permian "ophiuroid arm-ossicles" figured plate 6, figs. 1a-c, and figs 2a,b.] [see Imbrie 1955]
- Kozłowski, Roman. 1923. Faune Devonienne de Bolivie. *Ann. de Paleont.*, t. XII, 112 pp., X pls. [*Aspidosoma?* sp.; *Echinasterella?* sp.]
- Kozur, H. and H. Mostler. 1989. Echinoderm remains from the Middle Permian (Wordian) from Sosio Valley (Western Sicily). *Jb. Geol. B.-A.* 132(4):677-685.
- Kroh, A. 2003. First record of gorgonocephalid ophiuroids (Echinodermata) from the Middle Miocene of the Central Paratethys. *Cainozoic Research* 2(1-2):143-156. [Kroh 2006 erratum to table 2 in *Cainozoic Research* 4(1-2):119. [the erratum provides the last two columns of the table 2] [table 2 summarizes arm branching on the suborder Euryalina -- includes *Eospondylus* and *Onychaster* as non-branching, and *Kentrospondylus* as single branching].
- Kroh, A. 2006. [see Kroh 2003]
- Kühl, G., C. Bartels, D.E.G. Briggs, and J. Rust. 2012. Fossilien im Hunsrück-Schiefer. Goldschneck im Quelle & Meyer Verlag GmbH & Co., Wiebeisheim, 120 pp. [*Asteroidea*, *Baliactis tuberatus*, *Bundenbachia benecke*, *Encrinaster*, *Euzonosoma*, *Furcaster*, *Helianthaster rhenanus*, *Hystrigaster horridus*, *Loriolaster mirabilis*, Ophiuroidea, *Palaeostella solida*, *Protasteracanthion primus*, Schlangensterne, Seesterne, *Urasterella asperula*] [p. 39 fig. *Eospondylus primigenius* with crinoids; also p. 111 fig. with arm fragmentation]
- Kühl, G., C. Bartels, D.E.G. Briggs, and J. Rust. 2012. Visions of a vanished world: the extraordinary fossils of the Hunsrück Slate. [English edition of *Fossilien im Hunsrück-Schiefer*, with revisions]. Yale University Press, 128 pp. [p. 21 and Fig. 10 -- sponge-*Furcaster* association] [Fig. 36 ophiuroid stranded on *Thallocrinus* sp.] [Fig. 39 *Eospondylus* on crinoids *Haplocrinus* and *Triacrinus*] [pp. 80-93 stelleroids] [many significant comments on behavior and preservation] [p. 117, Figs. 119 *Eospondylus*, Fig. 120 *Urasterella*, autotomy]

May 11, 2016

- Kuhn, O. 1933. Die unterdevonischen Schiefer von Bundenbach. *Natur u. Kultur* 30:127-130, 6 figs. Innsbruck-Wien-München. [source Kutscher 1976] [not seen] [Aspidosoma]
- Kuhn, O. 1949. Lehrbuch der Paläozologie. 326 pp., 224 figs., 50 plates. Stuttgart. [source Kutscher 1976] [not seen] [*Euzonosoma tischbeinianum*, *Encrinaster roemeri*, *Helianthaster*]
- Kuhn, O. 1961. Die Tierwelt der Bundenbacher Schiefer. Neue Brehm-Bücherei, 274, 48 pp, 45 figs. Wittenberg-Lutherstadt. [source Kutscher 1976] [not seen] [figs. of *Encrinaster tischbeinianum*, *Roemeraster asperula*, *Furcaster palaeozoicus*, *Encrinaster roemeri*]
- Kukalová-Peck, J. 1973. A phylogenetic tree of the Animal Kingdom (including Orders and higher categories). *Nat. Mus. Canada, Publ. Zool. No. 8*, 78 pp. [source Sepkoski 1982]
- Kumpan, T. 2013. First find of the skeletal ossicles of eleutherozoid echinoderms from the Lišeň Formation (Famennian-Tournaisian) of the Moravian Karst. *Geologické výzkumy na Moravě a ve Slezsku*, Brno: Masarykova univerzita, 2013: 132-135. ISSN 1212-6209 [Furcaster, ?Eospondylus, stenurid sp.] [mainly vertebrae]
- Kumpan, T. 2013. Some Latest Famennian and Early Tournasian ophiuroids and holothuroids (Echinodermata) from the Moravian Karst. p. 32 in 14th Czech – Slovvak – Polish Palaeontological Conference, and 9th Polish Micropalaeontological Workshop, Abstract Volume, Grzybowski Foundation Special Publication No. 19, 2013. [Furcaster, ?Eospondylus, stenurid sp.] [mainly vertebrae]
- Kutscher, F. 1931. Zur Entstehung des Hunsrückschiefers am Mittelrhein und auf dem Hunsrück. *Jb. nass. Ver. Naturk.* 81:177-232, 2 figs, 2 plates. Wiesbaden. [source Kutscher 1976] [not seen] [Helianthaster rhenanus, Furcaster palaeozoicus, Encrinaster tischbeinianus, Urasterella asperula]
- Kutscher, F. 1963. Beiträge zur Sedimentation und Fossilführung des Hunsrückschiefers. 3. Die Anwendung der Röntgentechnik zur Diagnostik der Hunsrückschieferfossilien. *Notizbl. hess. L.-Amt Bodenforsch.* 91:74-86, 3 plates. Wiesbaden. [source Kutscher 1976] [not seen] [Urasterella asperula, Palaeostella solida, Medusaster rhenanus, Palaeonectria devonica]
- Kutscher, F. 1967. Die museale Darstellung von Hunsrückschiefer-Fossilien mit modernen Präparationsmethoden und Röntgendiagnostik. *Museumkunde*: 66-75, 4 figs. Berlin. [source Kutscher 1976] [not seen] [Urasterella verruculosa, Ophiurina lymani, Palaeactis lanceolatus, Palasterina tilmanni]
- Kutscher, Fritz. 1969. Beiträge zur Sedimentation und Fossilführung des

May 11, 2016

- Hunsrückschiefer-Fossilien. 23. Aus der Frühgeschichte der Untersuchung von Hunsrückschiefer-Fossilien. (Contributions to the sedimentation and fossil formation of the Hunsruck shales. 23. From the early history of the investigation of Hunsruck shale fossils.) *Dechiana* 122(1): 15-20, 1 pl. [English and French summaries][Euzonosoma tischbeinianum pl. i, figs. 5,6] [Urasterella asperula pl. i, figs. 2-4]
- Kutscher, F. 1970. Beiträge zur Sedimentation und Fossilführung des Hunsrückschiefers. 30. Die Echinodermen des Hunsrückschiefer-Meer. *Abh. hess. Landesamt. Bodenforsch.* 56:37-48, 2 figs., 1 pl. [A mass occurrence of Encrinaster roemeri described and figured] [also other topics such as water currents and arm layout]
- Kutscher, F. 1970. Die Vertseinerungen des Hunsrückschiefers. Erinnerungen an Walther Maximilian Lehmann. *Aufschluss*, 19. Sonderheft, Idar-Oberstein, pp. 87-100, 14 figs. Heidelberg. [source Kutscher 1976] [not seen] [Cheiropteraster giganteus, Urasterella verruculosa, Ophiurina lymani]
- Kutscher, F. 1970. Vor 350 Millionen Jahren. Leben im Hunsrückachiefer-See. *Mappe mit 16 Bildern und Beschreibungen.* Bad Kreuznach (Jung & Co.). [source Kutscher 1976] [not seen] [Euzonosoma tischbeinianum, Furcaster palaeozoicus, Verkiester 32-armed seastar, Medusaster rhenanus]
- Kutscher, F. 1970. Peter Tischbein und die Hunsrückschiefer-Fossilien. *Mitt. Ver. Heimatkd. Landkreis Birkenfeld u. Heimatfreunde Oberstein* 33:84-87, 2 figs. Idar-Oberstein. [source Kutscher 1976] [not seen] [Euzonosoma tischbeinianum]
- Kutscher, F. 1976. Beiträge zur Sedimentation und Fossilführung des Hunsrückschiefers. 43. Die Asterozoen des Hunsrückschiefers. -- *Geol. Jb. Hessen* 104:25-37. [complete review of all papers from Roemer 1862-1864 to 1972]
- Kutscher, F. and M. Horn. 1963. Beiträge zur Sedimentation und Fossilführung des Hunsrückschiefers. 4. Fossilien und organogener Detritus im Sediment. *Notizbl. hess. L.-Amt Bodenforsch* 91:87-91, 1 plate. Wiesbaden. [source Kutscher 1976] [not seen] [Palaeactis lanceolatus, Palasterina tilmanni, Miospondylus rhenanus, Ophiurina lymani, Furcaster palaeozoicus]
- Kutscher, Manfred. 2004. A sea-star (Echinodermata) in a Silurian Geschiebe (glacial erratic boulder) from Niederlehme near Berlin (Brandenburg). *Ueber einen Seestern (Echinodermata) in einem Silurgeschiebe von Niederlehme bei Berlin (Brandenburg).* *Archiv fuer Geschiebekunde* 3(8-12):619-626. [in German with English summary] [Urasterella ruthveni, U. ruthveni leintwardensis] [ZR2004] [complete review of all occurrences] [good photos; cross-section; isolated adams; body fossils]
- Kutscher, M. & L. Villier. 2003. Ophiuroid remains from the Toarcian of Sainte-Verge (Deux-

May 11, 2016

Sèvres, France): paleobiological perspectives. *Geobios* 36:179-194. [title also in French]
[Late Permian mass extinction and Early Triassic to Middle Jurassic burst of lineages sets
apart the Paleozoic and post-Paleozoic ophiuroid faunas] [excellent paper]